

Phenomenological parametrization of the charged particle
multiplicity distributions in restricted rapidity intervals in
 e^+e^- annihilation into hadrons and e^+p scattering at
HERA¹

O. G. TCHIKILEV²

*Institute for High Energy Physics
142284, Protvino, Russia*

Abstract

It is shown that charged particle multiplicity distributions in restricted (pseudo)rapidity intervals in e^+e^- annihilation and in e^+p scattering at HERA are quite well described by the modified negative binomial distribution and its simple extension.

(to be published in Phys. Lett. B)

¹This work was supported in part by INTAS, contract INTAS-93-3602

² E-mail: tchikilov@mx.ihep.su

Recently it has been shown [1-5] that negatively charged particle multiplicity distributions in e^+e^- annihilation are well described by the modified negative binomial distribution (MNBD) with the parameter Δ being practically energy independent and the parameter k approaching the value 7. It has been demonstrated also that the MNBD fairly well describes multiplicity distributions in lepton-nucleon interactions with a similar behaviour of the parameters [6]. The aim of this paper is to show that the MNBD and its simple extension describe quite well the charged particle multiplicity distributions in restricted rapidity intervals in e^+e^- annihilation into hadrons [7-12], and charged particle multiplicity distributions in restricted pseudorapidity intervals in e^+p scattering at HERA energies [13].

The MNBD is defined by the probability generating function

$$M(x) = \sum_n P_n x^n = \left(\frac{1 + \Delta(1-x)}{1 + r(1-x)} \right)^k, \quad (1)$$

where P_n is the probability to produce n particles. The MNBD can be considered as a convolution of the Newton binomial (the numerator in (1)) and the negative binomial (the denominator in (1)). The probabilities P_n for the MNBD can be computed also using iteration relations given in [4] or the formulae given in [14]. We have checked that all three methods give the same P_n values within the computer precision limits.

It has been assumed in [1,3,4] that the parameter k is the number of sources of particle production at some initial stage of the interaction; these sources develop independently of each other according to some branching process, producing intermediate neutral clusters. In this model the parameter $-\Delta$ is equal to the cluster decay probability into a charged hadron pair ($1 + \Delta$ is the probability of cluster decay into pair of neutral hadrons). One can associate these sources with the widely discussed clans [15] or superclusters [16]. A priori it is not evident that the same distribution should be valid for restricted rapidity intervals; indeed the initial sources can move with respect to each other and therefore contribute differently to the given rapidity interval, the same can happen for the intermediate neutral clusters. Let us assume that a fixed number of initial sources (less than or equal to k) contributes to a given rapidity interval (y_1, y_2) , and both charged decay products of each neutral cluster hit the rapidity interval with the probability ϵ_2 and only one charged particle hits the interval with the probability ϵ_1 . The probabilities ϵ_1 and ϵ_2 are assumed to be the same for all neutral clusters, whatever their source. In this case the probability generating

function for the charged particle multiplicity distribution has the form (1), where x is replaced by

$$\varphi_2(x) = 1 - \epsilon_1(1 - x) - \epsilon_2(1 - x^2). \quad (2)$$

The probability ϵ_1 should increase with the size of the rapidity interval up to a maximum, and then go down to zero; whereas the probability ϵ_2 should increase monotonically from zero to one. One expects that this rapidity dependence can naturally explain odd-even fluctuations for rapidity intervals covering nearly the whole phase space. One expects also that the parameter k increases from some initial value (may be 1) to the maximum, observed over the whole phase space. For some rapidity intervals, when contributions from configurations with different number of sources are equally important, the above assumptions seem to be crude; in this case one can use a sum of distributions with different k values³.

Analytical formulae for P_n can be obtained using the decomposition of the quadratic term in the denominator of the probability generating function into the product of two linear terms. For $k = 1$ it leads to the following formulae for the probabilities:

$$P_0 = \frac{1 + \Delta(\epsilon_1 + \epsilon_2)}{1 + r(\epsilon_1 + \epsilon_2)}, \quad (3)$$

$$P_i = \frac{(1 - \Delta/r)(\beta_1^{(i+1)} - \beta_2^{(i+1)})}{(1 + r(\epsilon_1 + \epsilon_2))(\beta_1 - \beta_2)}, \quad i > 0, \quad (4)$$

where

$$\beta_1 = \frac{b_1}{2} + \sqrt{\frac{b_1^2}{4} + b_2}, \quad \beta_2 = \frac{b_1}{2} - \sqrt{\frac{b_1^2}{4} + b_2} \quad (5)$$

and

$$b_1 = \frac{r\epsilon_1}{1 + r(\epsilon_1 + \epsilon_2)}, \quad b_2 = \frac{r\epsilon_2}{1 + r(\epsilon_1 + \epsilon_2)}. \quad (6)$$

For k greater than one we have calculated the P_n as a k -th convolution of the multiplicity distribution with $k = 1$, given by the formulae (3) and (4). It has been checked that for $\epsilon_2 = 0$ these calculations give the same results as the MNBD formulae.

Probabilities P_n can be computed also using the recurrence relations for Gegenbauer polynomials $C_n^\lambda(t)$ with the generating function

$$\sum_{n=0}^{\infty} C_n^\lambda(t) z^n = (1 - tz + z^2)^{-\lambda} \quad (7)$$

³This approach, for example, is used in [17], where multiplicity distributions are parametrized by a weighted superposition of two negative binomials.

(see for example [18]). The case $\Delta = 0$ gives

$$P_{n+1} = \frac{n+k}{n+1} b_1 P_n + \frac{n+2k-1}{n+1} b_2 P_{n-1} \quad (8)$$

with b_1 and b_2 given by (6) and with initial conditions

$$P_0 = (1 + r(\epsilon_1 + \epsilon_2))^{-k}, \quad P_1 = k b_1 P_0. \quad (9)$$

These recurrence relations can be used also for the case with $r = 0$ if one replaces k by $-k$ and r by Δ , and the final probabilities can be calculated by the convolution of these two distributions with $\Delta = 0$ and $r = 0$. We have checked that this method gives the same results as the method based on formula (4).

The experimental data have been parametrized by the MNBD ($\epsilon_2 = 0$) and its extension with $\epsilon_2 \neq 0$. The parameter Δ was fixed at the value $\Delta = -0.76$ taken from fits of the e^+e^- data [4,5] and the parameter r was calculated from the mean charged multiplicity $\langle n \rangle$ using the relation

$$r = \Delta + \frac{\langle n \rangle}{k(\epsilon_1 + 2\epsilon_2)}. \quad (10)$$

The integer parameter k has been tested in the interval from 1 to 9, and the parameters ϵ_1 and ϵ_2 have been assumed to be nonnegative. One should note that the parameter Δ cannot be obtained directly from fits, indeed the same P_n can be obtained for other Δ values by simple rescaling of the ϵ_1 , ϵ_2 and r .

Tables 1 and 2 give the results of the fits to the charged particle multiplicity distributions in the symmetric rapidity intervals $|y| < y_{cut}$ for e^+e^- annihilation for the TASSO [8], DELPHI [9] and ALEPH [11] data⁴. In some cases when the results with $\epsilon_2 \neq 0$ are not given, the fits have stopped on the boundary $\epsilon_2 = 0$.

One can see from tables 1 and 2 that the quality of the fits is quite good for all data excluding the DELPHI measurement [9]. It is necessary to note that the χ^2/NDF values for the fits should be considered just indicative, since the full covariance matrix is not published and therefore the proper treatment of the correlations between measurements of the neighbour multiplicities is not possible⁵. Nevertheless, for the TASSO measurements one can compare these values with the χ^2/NDF values obtained for negative binomial fits [8] made to

⁴ The P_n for the TASSO measurements [8] were calculated from the KNO scaling function $\psi(z)$, compiled in the COMPAS data base, IHEP, Protvino.

⁵Typical uncertainties (r.m.s.) in measured multiplicities in recent high energy experiments range from ~ 2 at $n_{ch} = 8$ to ~ 4 at $n_{ch} = 30$ [11].

the unfolded distributions. The χ^2/NDF ratios for MNBD fits in general are smaller than for negative binomial fits. Especially it is true for the statistically most significant TASSO measurement at $\sqrt{s} = 34.8$ GeV. The parameter k tends to rise with increasing size of the rapidity interval. The parameters ϵ_1 and ϵ_2 in general follow the expected dependence on the size of the rapidity interval.

The big χ^2/NDF ratios for the DELPHI measurements [9] are explained by the shoulder structure caused by the superposition of events with different number of jets (for the discussion see [17,19]). This shoulder structure, well described by the Monte Carlo program JETSET, is parametrized in [17] by a weighted superposition of two negative binomials. Following this approach we have parametrized multiplicity distributions in restricted rapidity intervals by the weighted superposition of two modified negative binomials. Let us denote the probabilities P_n for MNBD with the form (2), and $\epsilon_2 = 0$ by $p_n(k, \Delta, r_1, \epsilon_1)$. Then the weighted sum is given by

$$P_n = \alpha p_n(k, \Delta, r_1, \epsilon_1) + (1 - \alpha) p_n(k + 1, \Delta, r_2, \epsilon_1), \quad (11)$$

where α is the probability for the contribution of the k sources with the parameter r_1 , whereas $1 - \alpha$ is the probability for the contribution of the $k + 1$ sources with the parameter r_2 . The results of fitting the multiplicity distributions measured by DELPHI [9] in symmetric rapidity intervals by the weighted sum (11) are given in table 3. One can see the χ^2/NDF ratios are close to one; nevertheless it is necessary to note that the fits have four free parameters. We have tried also to fit by this parametrization the multiplicity distributions [9] in asymmetric rapidity intervals $0 < y < y_{cut}$ (or $-y_{cut} < y < 0$). The results are given in table 4. The χ^2/NDF ratios are near 1 for all intervals, excluding the intervals with $y_{cut} = 1$ and $y_{cut} = 1.5$: for these intervals, the ratios are near 2.

The results of the fits of the MNBD, and its extension with $\epsilon_2 \neq 0$, to charged particle multiplicity distributions in the single hemisphere for e^+e^- annihilation into hadrons [7-10,12] are given in table 5. The χ^2/NDF ratios are of order 1 for practically all measurements. One should note also that the fits to the DELPHI measurements [12] for $b\bar{b}$ events and all flavours have smaller χ^2/NDF ratios than the “shifted” negative binomial fits given in [19]⁶. For example, the MNBD fit for the $b\bar{b}$ sample gives $\chi^2/NDF = 13.3/27$, and the fit with $\epsilon_2 \neq 0$ gives $\chi^2/NDF = 7.9/26$, whereas the best negative binomial fit with a shift in multiplicity by 2 units has $\chi^2/NDF = 20/26$ [19].

It is of interest to note that the k values for single hemispheres are close to the parameter k for MNBD fits to the multiplicity distributions in the whole

⁶The shifts in multiplicity are due to the decays of the heavy b -quarks.

phase space [4,5], whereas the parameter ϵ_1 is rather close to the value 0.5. The second observation can be explained if one assumes that the intermediate neutral clusters hit one or other hemisphere with equal probability.

The results of the fits to the H1 data [13] in the pseudorapidity intervals $1 < \eta < \eta_{cut}$ are given in table 6. The χ^2/NDF values are in general close to 1 both for $\epsilon_2 = 0$ and for $\epsilon_2 \neq 0$. It is of interest to note that the fits with $\epsilon_2 \neq 0$ have smaller χ^2/NDF than the MNBD fits mainly for the energy W above 150 GeV.

In conclusion, it is shown that the MNBD and its simple extension describe multiplicity data in the restricted (pseudo)rapidity intervals for e^+e^- annihilation and e^+p deep inelastic scattering at HERA quite well. The DELPHI measurements [9] in symmetric and asymmetric rapidity intervals are fairly well parametrized by the weighted superposition of two modified negative binomial distributions with the parameters k and $k+1$. The dependence of the parameters on the size of the rapidity interval is explained by simple assumptions, extending the model proposed in [1,3,4]. Further checks of this phenomenological model can be done, for example, by studying the multiplicity distributions for particles of the same charge.

Acknowledgements

I am indebted to A. De Angelis and W. Venus for reading the manuscript and for the suggested improvements. The use of the COMPAS data base is acknowledged.

References

- [1] P. V. Chliapnikov and O. G. Tchikilev, Phys. Lett. **B242** (1990) 275.
- [2] N. Suzuki, M. Biyajima and G. Wilk, Phys. Lett. **B268** (1992) 447.
- [3] P. V. Chliapnikov and O. G. Tchikilev, Phys. Lett. **B282** (1992) 471.
- [4] P. V. Chliapnikov, O. G. Tchikilev and V. A. Uvarov, Phys. Lett. **B352** (1995) 461.
- [5] O. G. Tchikilev, Phys. Lett. **B382** (1996) 296.
- [6] O. G. Tchikilev, “Modified negative binomial description of the multiplicity distributions in lepton-nucleon scattering”, to be published in Phys. Lett. B.
- [7] HRS Collab., M. Derrick et al., Phys. Rev. **D34** (1986) 34
- [8] TASSO Collab., W. Braunschweig et al., Z. Phys. **C45** (1989) 193.
- [9] DELPHI Collab., P. Abreu et al., Z. Phys. **C52** (1991) 271.
- [10] OPAL Collab., P.D. Acton et al., Z. Phys. **C53** (1992) 539.
- [11] ALEPH Collab., D. Busculic et al., Z. Phys. **C69** (1995) 15.
- [12] DELPHI Collab., P. Abreu et al., Phys. Lett. **B347** (1995) 447.
- [13] H1 Collab., S. Aid et al., “Charged particle multiplicities in deep inelastic scattering at HERA”, preprint DESY 96-160 and hep-ex/9608011, 1996.
- [14] N. Suzuki, M. Biyajima and N. Nakajima, Phys. Rev. **D53** (1996) 3582.
- [15] A. Giovannini and L. Van Hove, Z. Phys. **C30** (1986) 391.
- [16] C. C. Shih and P. Carruthers, Phys. Rev. **D34** (1986) 2710.
- [17] A. Giovannini, S. Lupia and R. Ugoccioni, Phys. Lett. **B374** (1976) 231.
- [18] Higher transcendental functions, vol.2, p. 174, Ed. A. Erdelyi, McGraw Hill, New York, 1953.
- [19] A. Giovannini, S. Lupia and R. Ugoccioni, “The negative binomial distribution in quark jets with fixed flavour”, preprint DFTT 50/96, MPI-PhT/96-80, LU TP 96-23 and hep-ph/9609306, August 28th, 1996.

Table 1: Results of the fits with fixed parameter $\Delta = -0.76$ to the charged particle multiplicity distributions measured by TASSO [8] as a function of the centre of mass energy $\sqrt{s}$ in restricted rapidity intervals $|y| < y_{cut}$. The last column gives the χ^2/NDF for the negative binomial fits taken from [8].

$\sqrt{s}$ (GeV)	y_{cut}	k	ϵ_1	ϵ_2	χ^2/NDF	$\chi^2/NDF(NB)$
14.0	0.5	3	0.151 $\pm$ 0.022	0	12.4/14	15/13
		2	0.391 $\pm$ 0.017	0.106 $\pm$ 0.029	9.7/13	
	1.0	5	0.108 $\pm$ 0.023	0	37.2/20	27/19
		7	0.164 $\pm$ 0.023	0	36.0/23	33/22
	1.5	5	0.372 $\pm$ 0.045	0.170 $\pm$ 0.061	36.8/21	
		5	0.372 $\pm$ 0.045	0.170 $\pm$ 0.061	36.8/21	
22.0	0.5	9	0.386 $\pm$ 0.023	0	85.0/25	75/24
		5	0.239 $\pm$ 0.021	0.544 $\pm$ 0.018	72.8/24	
	1.0	3	0.181 $\pm$ 0.022	0	3.0/14	4/13
		2	0.411 $\pm$ 0.020	0.119 $\pm$ 0.030	4.8/13	
	1.5	3	0.421 $\pm$ 0.027	0	4.6/21	19/20
		2	0.470 $\pm$ 0.045	0.379 $\pm$ 0.043	11.7/20	
34.8	0.5	5	0.283 $\pm$ 0.028	0	6.5/26	8/25
		5	0.337 $\pm$ 0.030	0.054 $\pm$ 0.91	6.4/25	
	1.0	9	0.186 $\pm$ 0.025	0	14.5/28	13/27
		8	0.269 $\pm$ 0.057	0.190 $\pm$ 0.076	11.9/27	
	1.5	3	0.153 $\pm$ 0.011	0	10.1/15	22/14
		2	0.404 $\pm$ 0.008	0.109 $\pm$ 0.015	6.9/14	
43.6	0.5	3	0.387 $\pm$ 0.013	0	49.3/26	102/25
		2	0.462 $\pm$ 0.018	0.403 $\pm$ 0.017	50.3/25	
	1.0	4	0.415 $\pm$ 0.015	0	34.7/31	74/30
		3	0.406 $\pm$ 0.034	0.413 $\pm$ 0.029	30.7/29	
	1.5	8	0.082 $\pm$ 0.015	0	28.1/34	27/33
		6	0.308 $\pm$ 0.039	0.204 $\pm$ 0.048	30.7/33	
43.6	0.5	3	0.080 $\pm$ 0.022	0	6.7/17	8/16
		2	0.377 $\pm$ 0.015	0.077 $\pm$ 0.034	7.1/16	
	1.0	3	0.315 $\pm$ 0.026	0	25.8/27	44/26
		2	0.451 $\pm$ 0.035	0.399 $\pm$ 0.034	21.4/26	
	1.5	5	0.105 $\pm$ 0.027	0	45.6/32	39/31
		7	0.117 $\pm$ 0.027	0	43.6/34	39/33

Table 2: Results of the fits with fixed parameter $\Delta = -0.76$ to the charged particle multiplicity distributions measured by DELPHI [9] and ALEPH [11] at $\sqrt{s} = 91.2$ GeV in the symmetric rapidity intervals $|y| < y_{cut}$.

Collaboration	y_{cut}	k	ϵ_1	ϵ_2	χ^2/NDF
DELPHI	0.5	2	0.155 ± 0.018	0	55.8/21
	1.0	2	0.631 ± 0.016	0	111.3/35
	1.5	3	0.481 ± 0.014	0	279.4/44
		2	0.362 ± 0.019	0.730 ± 0.018	295.6/43
	2.0	4	0.425 ± 0.014	0	222.5/48
		3	0.346 ± 0.030	0.606 ± 0.027	218.9/47
ALEPH	0.5	2	0.305 ± 0.057	0	5.1/19
	1.0	2	0.753 ± 0.068	0	11.6/33
		2	0.552 ± 0.145	0.301 ± 0.162	10.8/32
	1.5	3	0.519 ± 0.056	0	32.7/39
		2	0.418 ± 0.089	0.696 ± 0.082	16.7/38
	2.0	3	0.929 ± 0.054	0	10.5/45
		2	0.335 ± 0.066	0.981 ± 0.057	37.0/44

Table 3: Results of the fits by the superposition of the two weighted MNBDs to the charged particle multiplicity distributions measured by DELPHI [9] at $\sqrt{s} = 91.2$ GeV in restricted rapidity intervals $|y| < y_{cut}$.

y_{cut}	k	α	ϵ_1	r_1	r_2	χ^2/NDF
0.5	5	0.735 ± 0.022	0.058 ± 0.034	6.87 ± 4.62	15.28 ± 9.36	11.0/18
1.0	5	0.727 ± 0.013	0.140 ± 0.024	6.08 ± 1.27	12.31 ± 2.29	27.9/32
1.5	8	0.681 ± 0.010	0.062 ± 0.022	6.50 ± 1.24	13.03 ± 2.22	51.1/41
2.0	9	0.570 ± 0.011	0.131 ± 0.024	7.05 ± 1.47	13.19 ± 24.92	46.7/45

Table 4: Results of the fits by the superposition of the two weighted MNBDs to the charged particle multiplicity distributions measured by DELPHI [9] at $\sqrt{s} = 91.2$ GeV in restricted rapidity intervals $0 < y < y_{cut}$.

y_{cut}	k	α	ϵ_1	r_1	r_2	χ^2/NDF
0.5	4	0.790 ± 0.025	0.060 ± 0.039	3.96 ± 2.24	9.58 ± 6.78	3.6/12
1.0	4	0.848 ± 0.009	0.061 ± 0.029	9.50 ± 4.52	23.83 ± 11.38	35.4/25
1.5	6	0.769 ± 0.008	0.065 ± 0.022	8.05 ± 2.98	20.51 ± 7.03	55.8/29
2.0	9	0.685 ± 0.008	0.055 ± 0.021	7.84 ± 3.27	19.51 ± 7.48	75.5/30
3.0	9	0.423 ± 0.017	0.219 ± 0.027	2.24 ± 0.41	4.51 ± 0.62	36.5/30
5.0	9	0.563 ± 0.061	0.467 ± 0.025	1.44 ± 0.19	2.01 ± 0.16	31.5/30

Table 5: Results of the fits with fixed parameter $\Delta = -0.76$ to the charged particle multiplicity distributions in a single hemisphere for e^+e^- annihilation into hadrons.

Experiment	$\sqrt{s}$ (GeV)	k	ϵ_1	ϵ_2	χ^2/NDF
TASSO [8]	14.0	6	0.566 ± 0.009	0	10.5/17
		4	0.568 ± 0.027	0.292 ± 0.024	7.1/16
	22.0	8	0.456 ± 0.010	0	5.5/17
		6	0.535 ± 0.024	0.162 ± 0.030	5.0/16
	34.8	8	0.470 ± 0.005	0	26.4/23
		6	0.541 ± 0.015	0.196 ± 0.017	24.7/22
	43.6	9	0.376 ± 0.010	0	10.6/23
		6	0.483 ± 0.033	0.253 ± 0.033	12.5/22
HRS [7]	29.0	9	0.487 ± 0.011	0	17.8/18
		5	0.576 ± 0.047	0.321 ± 0.039	24.5/17
OPAL [10] with K_s, Λ	91.2	5	0.653 ± 0.018	0	18.7/34
		5	0.667 ± 0.080	0.245 ± 0.078	12.1/33
OPAL [10] no K_s, Λ	91.2	5	0.782 ± 0.018	0	7.6/32
		5	0.684 ± 0.086	0.150 ± 0.100	7.0/31
DELPHI [9]	91.2	6	0.671 ± 0.008	0	98.7/34
		4	0.309 ± 0.023	0.729 ± 0.018	80.7/33
DELPHI [12], $b\bar{b}$ -events	91.2	6	0.844 ± 0.011	0	13.3/27
		5	0.592 ± 0.068	0.452 ± 0.062	7.9/26
DELPHI [12], all flavours	91.2	6	0.691 ± 0.012	0	24.9/29
		5	0.506 ± 0.057	0.412 ± 0.048	20.1/28

Table 6: Results of the fits with fixed parameter $\Delta = -0.76$ to the charged particle multiplicity distributions measured by H1 [13] in restricted pseudorapidity intervals $1 < \eta < \eta_{cut}$.

W(GeV)	η interval	k	ϵ_1	ϵ_2	χ^2/NDF
80 $\div$ 115	1 $\div$ 2	3	0.038 $\pm$ 0.073	0	0.8/13
		1	0.547 $\pm$ 0.015	0.303 $\pm$ 0.047	14.3/12
	1 $\div$ 3	4	0.100 $\pm$ 0.076	0	1.8/17
		3	0.339 $\pm$ 0.102	0.167 $\pm$ 0.134	2.2/16
	1 $\div$ 4	9	0.026 $\pm$ 0.079	0	0.5/18
		5	0.317 $\pm$ 0.215	0.229 $\pm$ 0.228	0.7/17
	1 $\div$ 5	9	0.162 $\pm$ 0.044	0	1.6/18
		3	0.346 $\pm$ 0.056	0.627 $\pm$ 0.060	9.9/17
115 $\div$ 150	1 $\div$ 2	2	0.199 $\pm$ 0.035	0	6.9/14
		1	0.533 $\pm$ 0.026	0.285 $\pm$ 0.027	62.5/13
	1 $\div$ 3	3	0.178 $\pm$ 0.090	0	2.8/18
		1	0.588 $\pm$ 0.060	0.385 $\pm$ 0.055	113.3/17
	1 $\div$ 4	7	0.043 $\pm$ 0.056	0	4.5/20
		7	0.174 $\pm$ 0.048	0.122 $\pm$ 0.110	3.4/19
	1 $\div$ 5	9	0.154 $\pm$ 0.047	0	7.6/21
		4	0.319 $\pm$ 0.058	0.515 $\pm$ 0.056	12.8/20
150 $\div$ 185	1 $\div$ 2	3	0.028 $\pm$ 0.092	0	2.9/14
		3	0.182 $\pm$ 0.055	0.066 $\pm$ 0.042	1.0/13
	1 $\div$ 3	3	0.195 $\pm$ 0.116	0	4.0/20
		2	0.442 $\pm$ 0.170	0.322 $\pm$ 0.097	3.0/21
	1 $\div$ 4	6	0.064 $\pm$ 0.083	0	4.4/22
		6	0.162 $\pm$ 0.083	0.178 $\pm$ 0.097	3.0/21
	1 $\div$ 5	9	0.088 $\pm$ 0.064	0	3.4/22
		9	0.218 $\pm$ 0.034	0.154 $\pm$ 0.088	1.5/21
185 $\div$ 220	1 $\div$ 2	3	0.034 $\pm$ 0.096	0	2.4/14
		3	0.173 $\pm$ 0.064	0.056 $\pm$ 0.041	0.6/13
	1 $\div$ 3	3	0.188 $\pm$ 0.096	0	3.8/21
		2	0.366 $\pm$ 0.075	0.322 $\pm$ 0.059	8.4/20
	1 $\div$ 4	5	0.126 $\pm$ 0.091	0	9.2/22
		4	0.232 $\pm$ 0.060	0.300 $\pm$ 0.067	8.1/21
	1 $\div$ 5	9	0.169 $\pm$ 0.074	0	8.7/23
		5	0.233 $\pm$ 0.060	0.502 $\pm$ 0.064	6.2/22